

International Journal of Modern Physics A
 © World Scientific Publishing Company

PRIMARY ENERGY MEASUREMENT WITH EAS CHERENKOV LIGHT: EXPERIMENT QUEST AND CORSIKA SIMULATION

E. KOROSTELEVA, L. KUZMICHEV, V. PROSIN
Skobel'tzyn Institute of Nuclear Physics, MSU, Moscow, Russia

B. LUBSANDORZHIEV
Institute of Nuclear Research, RAS, Moscow, Russia

EAS-TOP COLLABORATION
INFN, IFSI, Universita' Torino, Italy

Received (Day Month Year)
 Revised (Day Month Year)

A new primary mass independent method of energy measurement has been developed by exploiting: a) the joint analysis of the shower size, obtained by EAS-TOP, and of the EAS atmospheric Cherenkov light lateral distribution, obtained by the QUEST array, and b) simulations based on the CORSIKA/QGSJET code. The method is based on the correlation between the size/energy ratio and the steepness of Cherenkov light lateral distribution and has been compared with a "classical" one based on the Cherenkov light flux at a fixed distance (175 m) from the EAS core. An absolute energy calibration of the EAS atmospheric Cherenkov light flux has been obtained.

Keywords: Primary energy; EAS Cherenkov light.

1. Experiment QUEST and Simulations

The QUEST experiment was developed to combine atmospheric Cherenkov light data to the charged particle EAS-TOP measurements (Gran Sasso, Italy, 2000 m a.s.l.)¹ The Cherenkov light detector was based upon five wide angle QUASAR-370 (37 cm diameter) semispheric photomultipliers installed on five telescopes (average pointing at the direction $\theta = 34^\circ$, $\varphi = 167^\circ$, full field of view for $\theta \leq 40^\circ$, used in analysis, is 0.41 sr).

A new fitting function has been used to derive two main parameters of EAS Cherenkov light LDF for every recorded event: the light flux at core distance of 175 m Q_{175} and the LDF steepness, defined as the ratio of the fluxes at 100 and 200 m from the axis:¹ $P = Q(100)/Q(200)$.

The energy measurement methods are based on parameterizations of Cherenkov light lateral distributions obtained from simulations performed through CORSIKA/QGSJET²⁻⁴ for primary protons and iron nuclei with total energy 1, 2, 4 and 8 PeV, and zenith angles θ from 0° to 39° .

The experiment has been additionally simulated by using such parameterizations and


2 *E. Korosteleva et al.*

Fig. 1. CORSIKA: Correlation between EAS Cherenkov LDF steepness and N_e/E_0 .




Fig. 2. EXPERIMENT: Comparison of energy, obtained with two different methods for $E_{CHER} \geq 2 \times 10^{15}$ eV.

extracting the primary energy, the depth of EAS maximum $X_{\max}$, shower axis direction and coordinates. The primary composition is assumed 50% p and 50% Fe. All other parameters have been calculated as functions of the main (i.e. $P = F_1(X_{\max}, \theta, A)$, $N_e = F_2(E_0, P, A)$ and so on). All the known apparatus errors and fluctuations have been taken into account.

2. 1st Method: Size and Cherenkov Light LDF steepness P

To derive the size comparable to the experimental one, we have taken into account both electrons and muons and used the experimental procedure of size reconstruction with NKG fitting function. Figure 1 shows the correlation between the parameter P ($P = Q(100)/Q(200)$) and the ratio of the size to primary energy (N_e/E_0). Using this relation we can get the primary energy from measurement of N_e and P :

$$E_{SIZE} [\text{eV}] = 1.59 \times 10^{11} N_e / 10^{0.33P}.$$

This method is quite composition independent, since it gives only 6% of difference in energy estimation for proton and iron. Its main advantage relies in the well developed technique of scintillator response calibration based on the measurement of the single particle response.⁵ From the simulation we derive for the reconstructed energy relative errors (mostly due to apparatus error of parameter P reconstruction) of about 30%.

Similar method of energy reconstruction, but for LDF steepness, estimated at smaller distances from the core (20 – 100 m), was suggested in Ref. 6.

3. 2nd Method: Cherenkov Light Flux at 175 m Core Distance Q_{175} .

From the simulation we obtain the following relation between Q_{175} [photon $\cdot$ cm $^{-2}$ $\cdot$ eV $^{-1}$] and primary energy E_{CHER} [eV]:

$$E_{CHER} = 4 \times 10^{14} Q_{175}^{0.94}.$$

This method leads to larger differences in energy estimations for primary proton and iron (about 19%), than the first one. But the error of reconstructed energy (estimated with the above mentioned simulation of the experiment) is about 15% even for the assumed “50% p + 50% Fe” composition, i.e. two times better than the first one.

The main problem of this method is in the absolute calibration of Cherenkov light detectors, its error being estimated between 18% to 30% for different experiments. To get better accuracy we used the most probable ratio E_{SIZE}/E_{CHER} as the coefficient for absolute calibration of Cherenkov detectors response. The coefficient is estimated for the natural mass composition. After such recalculation, the final comparison of two methods in the experimental data is shown in Fig. 2. The standard deviation of the experimental distribution is very close to the one obtained in MC simulation of experiment for the SIZE/CLDF method (see above).

4. A Reference Integral Primary Cosmic Rays Intensity

600 events have been recorded during 140 h of data taking with energy (derived with the “re-calibrated Cherenkov light flux method”) larger than 3×10^{15} eV. The corresponding integral intensity is:

$$I(E_0 \geq 3 \times 10^{15} \text{ eV}) = (2.3 \pm 0.1) \times 10^{-7} \text{ m}^{-2} \cdot \text{s}^{-1} \cdot \text{ster}^{-1}.$$

The statistical uncertainty of the measurement is about 5%.

This estimation can be used as a reference point for other experiments having no precise absolute calibration, such as Cherenkov light experiment TUNKA.⁷

Acknowledgements

The experimental start of this work has been supported by INTAS grant 96-0526.

References

1. E. Korosteleva, L. Kuzmichev, V. Prosin, and EAS-TOP Collab. in *Proc. 28th ICRC*, Tsukuba, Japan, **1**, 89 (2003).
2. D. Heck *et al.*, FZKA report 6019 (1998).
3. N. N. Kalmykov *et al.*, *Nucl. Phys. B (Proc. Suppl.)* **52B**, 17 (1997).
4. S. Martinez *et al.*, *Nucl. Instrum. Meth.* **A357**, 567 (1995).
5. M. Aglietta *et al.* (EAS-TOP Collab.), *Nucl. Instrum. Meth.* **A336**, 310 (1993).
6. A. Lindner, *Astropart. Phys.* **8**, 235 (1998).
7. D. Chernov *et al.*, in *Proc. 27th ICRC*, Hamburg, Germany, **2**, 581 (2001).